

REPORT

STAT

DATE DISTR. 23 June 1948

NO. OF PAGES 1

PLACE ACQUIRED USSR

NO. OF ENCLS.
(LISTED BELOW)

DATE OF
INFORMATION 1947

SUPPLEMENT TO
REPORT NO.

STAT

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE OF THE UNITED STATES WITHIN THE MEANING OF THE ESPIONAGE ACT 50 U.S.C. 31 AND 32, AS AMENDED. ITS TRANSMISSION OR THE REVELATION OF ITS CONTENTS IS ANY WARNER TO AN UNAUTHORIZED PERSON IS PROHIBITED BY LAW. PROSECUTION OF THIS FORM IS PROHIBITED. HOWEVER, INFORMATION CONTAINED IN THIS FORM MAY BE UTILIZED AS DEEMED NECESSARY BY THE RECEIVING AGENCY.

THIS IS UNEVALUATED INFORMATION FOR THE RESEARCH
USE OF TRAINED INTELLIGENCE ANALYSTS

SOURCE Documentary as indicated. (Information specifically requested.)

RECENTLY PUBLISHED RESEARCH OF THE INSTITUTE OF
BIOLOGICAL PROPHYLAXIS OF INFECTIONS, USSR

"Specific Polysaccharide Complexes of Microorganisms,"
A. M. Kozin, I. S. Buyanovskaya, A. M. Rykalova, N. I. Kuzina, Inst Biol Prophylaxis of Infections, Moscow

"Biekhimiya" Vol 12, 1947, pp 340-3

Polysaccharide complexes from tissues of guinea pigs, white mice, and human subjects were isolated by modified known methods. Polysaccharide from guinea pigs (yield 1.0%) contained 4.5-6.8% N. After hydrolysis with 0.1N HCl, the mixture analyzed: polysaccharide 59%, lipides 9%, peptides 16%, water-soluble fraction 36%. Polysaccharide from human tissues (yield 0.2-0.4%) contained 2.5% N. Reducing substances after hydrolysis with 1 N H₂SO₄ for 4 hours varied from 11 to 26%. Polysaccharide from white mice (yield 0.9%) had 5.8% N and yielded 17.5% reducing substances (as glucose) on hydrolysis with 1 N H₂SO₄ for 4 hours. Chemical composition of polysaccharide was similar to that of the guinea pig. Glycogen was not a component of any polysaccharide. All three complex polysaccharides possessed specific antigenic properties.

- END -

- 1 -

CLASSIFICATION: RESTRICTED

STATE		NAVY		X		NSRB		DISTRIBUTION	
ARMY	X	AIR	X	RDB	X				

RESTRICTED